

ANTIDEPRESSANT-INDUCED SEXUAL DYSFUNCTION

Jon-Paul Khoo

Objectives

- To appreciate the relationship between major depressive disorder, its treatment and sexual dysfunction
- To review the assessment of sexual function
- An approach to the clinical management of antidepressant-induced sexual dysfunction

Introduction

- MDD is a critical public health problem
- Antidepressant use is widespread
 - ▣ 264 million prescriptions filled for antidepressants in US in 2011, accounting for US\$11 billion (1)
 - ▣ Antidepressants were the 4th most commonly prescribed group of medications in 2009 (2)
- Sexual dysfunction is a common side-effect
- Sexual dysfunction is distressing, contributes to poor psychosocial functioning, and contributes to poor treatment compliance

1. Lindsley C. ASC Chem Neurosci 2012; 3 (8):630-1.
2. Lindsley C. ASC Chem Neurosci 2010; 1(6):407-8.

What is sexual dysfunction, how frequently does it occur and how do we evaluate it?

Sexual dysfunction (SD)

- Dysfunction in phases of the sexual response cycle
 - ▣ Desire: low/absent
 - ▣ Arousal: erectile dysfunction/reduced vaginal lubrication
 - ▣ Orgasm: delayed/anorgasmia; spontaneous; ejaculatory disorder
- Pain and sensory changes
- (What about too much sexual function?)

SD in the general population

US epidemiological data, 18-59yo

Women 30%

- 32% lack interest
- 26% inability to orgasm
- 23% absent pleasure
- 21% lubrication problems
- 15% dyspareunia
- 12% performance anxiety

Men 45%

- 31% premature ejaculation
- 18% performance anxiety
- 15% lack interest
- 10% erectile dysfunction
- 8% inability to orgasm
- 8% absent pleasure

Laumann E et al. JAMA 1999;281:537-544.
Nurnberg H. Drugs Today 2008;44(2):147-168.

Assessment of sexual functioning

- Assess premorbid/baseline sexual functioning
- Consider psychiatric and psychological issues that might contribute to sexual difficulties
- Consider physical health comorbidities and concomitant medication
- Consider alcohol and illicit substance usage
- Assess intra-morbid sexual functioning in current (and past) depressive episodes
- Current psychosocial context
- Personal importance of sexual activity and current relationship impact of SD

Sexual dysfunction in MDD


Sexual dysfunction and MDD

- In untreated depressed patients (MDD, dysthymia, recurrent brief depression), SD occurs in up to 50% (40-65%) vs 24% controls (1,2)
- MDD-induced SD (2)
 - Low interest: 40% men and >50% women
 - Arousal dysfunction 40-50%
 - Orgasm difficulty 15-20%
- Greater depressive severity, duration and recurrence predicts more SD (3,4)
- Antidepressant treatment improves SD in those with depression-induced SD (5,6): NB attribution bias

1. Angst J. *Int Clin Psychopharmacol*. 1998 Jul;13 Suppl 6:S1-4. 2. Kennedy S et al. *J Affect Disord*. 1999 Dec;56(2-3):201-8.
 3. Bonierbale M et al. *Curr Med Res Opin*. 2003;19(11):141-24. 4. Czarowski J et al. *Arch Sex Behav*. 2004;33:239-48.
 5. Baldwin D et al. *J Psychopharmacol*. 2006;20:91-6. 6. Baldwin D et al. *Hum Psychopharm Clin*. 2008;23:527-32.

The relationship between MDD and SD


Clayton A et al. *J Sex Med*. 2009 May;6(5):1200-11. Kennedy S et al. *J Clin Psychopharmacol*. 2009;29(2):157-64. Gregorian R et al. *Ann Pharmacother*. 2002;36(10):1577-89. Rosenberg K et al. *J Sex Marital Ther*. 2003;29(4):289-96.



Antidepressant-induced SD

Monoamine mechanisms of AISD

- Reduced desire
 - Serotonin reuptake blockade reduces mesolimbic dopaminergic neurotransmission via 5HT2 agonism
- Arousal dysfunction
 - SNS and PNS spinal reflex inhibition by serotonin
- Orgasm dysfunction
 - Reduced dopaminergic and noradrenergic neurotransmission by 5HT2 agonism



Clayton A et al. *Psychiatr Clin N Am* 2016;39:427-463.

Antidepressant-induced SD

- Worsening of pre-existing SD or new-onset SD
- Only 1/4 will disclose
 - ▣ Spontaneous disclosure 14% vs 60% on questionnaire
- AISD in women
 - ▣ 27-65%
 - ▣ More SD than men
 - ▣ More interest and orgasm dysfunction than men
 - ▣ Less likely to discuss SD and more likely to attribute SD to other causes
- AISD in men
 - ▣ 26-57%
 - ▣ More severe SD than women
 - ▣ More arousal dysfunction than women

Serretti A and Chessa A. J Clin Psychopharmacol 2009;29(3):259-266. Montejo A et al. J Clin Psychiatry 2001;62(suppl3):10-21. Williams V et al. J Clin Psychiatry 2006;67:204-10. Williams V et al. J Psychopharmacol 2010;24:489-96. Reichspoder U et al. Drug Safety 2014;37:119-31.

Consequences of AISD

- Non-adherence
 - ▣ SD is one of the most common side-effects leading to treatment discontinuation
 - ▣ 42% of men and 15% women discontinue antidepressants due to concerns about SD*
- Depressive relapse
- Reduced quality of life
- Tolerance for AISD reduces with increasing time in recovery


* Rosenberg K et al. J Sex Marital Ther 2003;29:289-296. van Geffen E et al. Eur J Clin Pharmacol. 2007;63:1193-1199. Gregorian K et al. Ann Pharmacother. 2002;36(10):1577-89. Clayton A et al. J Sex Med. 2009 May;6(5):1200-11. Reichspoder U et al. Drug Safety 2014;37:119-31.

AISD in antidepressant RCTs

- AISD occurs in 15-80%
- Mean total placebo rate of SD on MA = 14.2%
- Variation in incidence is multifactorial, and due to methodological and attributional difficulties
 - ▣ No standard SD definitions/classification
 - ▣ SD is rarely a primary or secondary outcome measure, often requiring spontaneous disclosure
 - ▣ Variable rating scale sensitivity
 - ▣ No baseline (premorbid/pre-treatment) levels reported
 - ▣ Doctors consistently underestimate SD in depressed patients
- Note: observational studies in naturalistic settings report higher rates of SD than RCTs

Serretti A and Chessa A. J Clin Psychopharmacol 2009;29(3):259-266. Montejo A et al. J Clin Psychiatry 2001;62(suppl3):10-21. Gartlehner G et al. Ann Intern Med 2011;155(11):772-85. La Torre A et al. Pharmacopsychiatry 2013;46:191-199. Balon R. Am J Psychiatry 2006;163:1504-9.

SD: attributable fractions



Meta-analysis of AISD

Serretti A and Chessa A. J Clin Psychopharmacol 2009;29(3):259-266.

SD occurs across all phases with all antidepressants, except where exceptions are noted
Mean total placebo rate of SD was 14.2%

>placebo		≤placebo
HIGHEST RISK	MIDDLE RISK	LOWEST RISK
Sertraline 80%	Imipramine 44%	Mirtazapine 24%**
Venlafaxine 80%	Phenelzine 42%	Bupropion 10%***
Citalopram 79%	Duloxetine 42%	Moclobemide 4%
Paroxetine 71%	Escitalopram 37%*	Agomelatine 4%
Fluoxetine 70%	Fluvoxamine 26%	Vortioxetine ^Δ

* Escitalopram has a placebo level of desire dysfunction

** Mirtazapine has only desire dysfunction

*** Bupropion has only arousal dysfunction

^Δ Not part of this study, but ~placebo (with and without baseline SD) a pooled analysis of 7 RCTs

Important observations regarding AISD

- Minimal clinical differences in AD efficacy, but differences in onset, adverse events (AISD) and rates of discontinuation (1)
- Intra- and inter-class variation
- Dose-related
- Usually occurs early in treatment
- Typically persists throughout treatment
- Typically resolves on discontinuation of the offending treatment

1. Gartlehner G et al. Ann Intern Med 2011;155(11):772-85.

The clinical management of AISD

Clinical Management of AISD

- Assessment of sexual functioning prior to treatment
- (A priori prescription of low SD risk treatment in those already suffering SD or very concerned about developing it)
- Validated tools for qualifying/quantifying SD

Validated depression-specific SD questionnaires

Scale	No. of Items	Method of administration	Time required (mins)	Gender Versions?	Comments
ASEX	5	Clinician	5-10	Yes	Simple design
CSFQ-short	14	Self	5	Yes	Has long version
PRSexDQ or SALSEX	7	Clinician	5	Yes	Specific to AISD
SexFX	11	Clinician/self	5-10	Yes	Specific to AISD

Direct enquiry

ASEX: Arizona Sexual Experiences Scale McGilvey C et al. J Sex Marital Ther 2000;26:25-40.
 CSFQ: Changes in Sexual Functioning Questionnaire Clayton A et al. Psychopharmacol Bull 1997;33:731-745.
 PRSexDQ: Psychotropic-related Sexual Dysfunction Questionnaire Montejo A et al. J Clin Psychiatry 2001;62:10-21.
 SexFX: Sex Effects Scale Kennedy S et al. J Clin Psychiatry 2000;61(4):276-281.

Clinical Management of AISD

- Assessment of sexual functioning prior to treatment
- (A priori prescription of low SD risk treatment in those already suffering SD or very concerned about developing it)
- Validated tools for qualifying/quantifying SD
- Rule out causes unrelated to depression or ADT
 - Physical/medication/substance/psychological/social
- Target baseline
- Explanation and education
- Specific strategies
 - Behavioural
 - Pharmacological
 - Complimentary

Behavioural management of AISD

- Exercise
 - 3/52 moderate strength training and aerobic exercise 30/60 before sexual activity improved sexual desire and function in women (1)
- Scheduling sexual activity
 - May increase orgasm function in women (2)
- Changing sexual technique
- Vibratory stimulation
- Psychotherapy

- There are no RCTs of behavioural strategies

1. Lorentz T et al. Am Behav Med 2012;43(3):352-61.
 2. Lorentz T et al. Depress Anxiety 2014;31(3):188-95.

Pharmacological management of AISD

- Watchful waiting
- Scheduling sexual activity
- Dose reduction
- Drug holiday
- Switch
- Augment

Watchful waiting

- Waiting for tolerance
 - ▣ Adaptation occurs in ~5-10% over 4-6/12 (1-3)
- Where patient is
 - ▣ Experiencing good antidepressant efficacy;
 - ▣ Considered to be on short-term treatment; and
 - ▣ Accepting AISD as a “price worth paying”
- Very ineffective: the vast majority will have no improvement over 6/12 (4) therefore such should not be expected

1. Ashton A et al. J Clin Psychiatry 1998;59(3):112-5. 2. Montejo A et al. J Clin Psychiatry 2001;61(suppl3):10-21.
3. Clayton A et al. J Clin Psychiatry 2006 (suppl6):33-7. 4. Montejo-Gonzalez A et al. J Sex marital Ther 1997;23(3):176-94.

Scheduling

- Schedule sexual activity at trough serum level
- Where
 - ▣ Antidepressant has short half-life (eg sertraline, paroxetine, clomipramine)
- Limited benefit for the majority and reduces spontaneity

Dose reduction

- AISD appears to be dose-related
- Lowest therapeutic treatment dose (“dose inflation”)
- Strategies usually recommend a 50% dose reduction
- Where
 - ▣ Antidepressant has a flat dose-response curve
 - ▣ Patient is experiencing good antidepressant efficacy
- Difficulties include reducing SD at the cost of symptom control, discontinuation and nonadherence

Drug holiday

- Temporary reduction or suspension of antidepressant
- Most methods use a brief abstinence strategy
 - ▣ E.g. cease after dose on Thursday- restart Sunday (1): improved sexual function 50% of the time on weekends without mood deterioration
- Where
 - ▣ Patient is experiencing good efficacy
 - ▣ Antidepressant has a shorter half-life
 - ▣ Sexual activity is relatively infrequent
- Overall, not very effective
- Difficulties include discontinuation, relapse, reduced spontaneity, conflicting messages about adherence and non-adherence
- ONLY recommend if in full remission and SD impairment is so severe that the sufferer would otherwise cease treatment


1. Montejo A et al. J Clin Psychiatry 2001;61(suppl3):10-21.

Switch


- Change antidepressant to a lower SD risk alternative
 - ▣ Agomelatine, vortioxetine, bupropion, moclobemide, mirtazapine
 - ▣ SSRI to SSRI probably wont work, though SSRI to SNRI might
- Where
 - ▣ Suboptimal efficacy
 - ▣ Treatment refusal due to SD
- High likelihood of alleviating AISD
- Difficulties include other side-effects and risks with new treatment, potential loss of efficacy, potential crossover
- Single RCT
 - ▣ Vortioxetine > Escitalopram in improving TESD in those taking SSRI (citalopram, paroxetine, sertraline) whilst maintaining efficacy (1)

1. Clayton A et al. Expert Opin Drug Saf 2014;13(10):1361-1374.


ADT switching strategies



- + Rapid, always medicated
- More interactions/side-effects



- + For partial responders so no efficacy loss
- More interactions/side-effects but easier to identify which from



- + Cleanest
- Takes longer (especially with washout), risks deterioration in crossover (therefore best for non-responders)

DPG 2015; Malhi G et al. ANZJP 2015; 49: 1087-1206.

Augmentation “antidote”

- Adding a second treatment to improve sexual functioning
 - Pulse vs regular?
- Where
 - Patient is experiencing good efficacy
 - Patient accepting additional medication
- Difficulties include drug-drug interactions; side-effect synergy, increased cost, the complexity of treatment regimen might reduce adherence, and, in Australia, none are indicated or reimbursed
- Some adjunctive therapies might also enhance antidepressant response (eg bupropion, mirtazapine, agomelatine)
- Limited database

Which ‘antidotes’?

- Cochrane database review 2103 (1)
 - PDE inhibitors
 - Sildenafil (3 studies (2-4))
 - Men and women reported improved sexual function and satisfaction
 - Most favoured strategy for men with erectile dysfunction
 - Women had reduced orgasm disturbance
 - Tadalafil (2 studies (5-6))
 - ED due to antidepressants
 - Bupropion
 - 150mg BD (3 studies (7-9)) > placebo with no deterioration in mood
 - 150mg OD (2 studies (10-11)) = placebo
 - ‘The most promising approach studies so far’ in women with AISD

1. Taylor M et al. Cochrane Database of Systematic Reviews 2013; 5. Art. No.: CD003382. 2. Numburg H et al. JAMA 2008;300(4):395-404.
 3. Numburg H et al. JAMA 2003;289(1):56-64. 4. Fava M et al. J Clin Psychiatry 2006;67(2):240-6. 5. Eshbaugh J et al. Urology 2011;75(5):1137-41.
 6. Segroves R et al. J Clin Psychopharmacol 2007;27(1):62-66. 7. Ollin M et al. J Sex Med 2002; 28: 131 - 138.
 8. Nordan M. Depression 1994; 12:109 - 112. 9. Safarinejad M. BJU Int 2010; 106: 840 - 847. 10. Safarinejad M. J Psychopharmacol 2011; 25: 370 - 378. 11. Thase M. J Clin Psychiatry 2005; 66: 1974 - 1981.

Other ‘antidotes’

- Controlled data
- Case reports/series
- Complimentary therapies with controlled data

Other ‘antidotes’

- Controlled data
- Case reports/series
- Complimentary therapies with controlled data
- Mirtazapine
- Aripiprazole
- Testosterone

Other ‘antidotes’

- Controlled data
- Case reports/series
- Complimentary therapies with controlled data
- Buspirone
- Cyproheptdine/loratadine
- Yohimbine
- Dopamine agonists: amantadine, bromocriptine and psychostimulants
- Anticholinergics

Other ‘antidotes’

- Controlled data
- Case reports/series
- Complimentary therapies with controlled data
- Maca root
- Saffron
- SAME

Take home messages

Objective 1: take home message

- **The relationship between major depressive disorder, its treatment and sexual dysfunction**
 - AISD includes both aggravations of existing SD and new onset SD
 - In the depressed individual on treatment who experiences SD, the cause may be due to the underlying depressive morbidity, to the treatment, or to some pre-morbid background physiological, illness, substance or psychosocial variable

Objective 2: take home message

- **The assessment of sexual function**
 - Premorbid/baseline sexual functioning
 - Psychiatric and psychological issues associated with SD
 - Physical health comorbidities and concomitant medication
 - Alcohol and illicit substance usage
 - Intra-morbid sexual functioning in current (and past) depressive episodes
 - Current psychosocial context
 - Personal importance of sexual activity and current relationship impact of SD

Objective 3: take home message

- **The clinical management of AISD**
 - Rule out causes unrelated to depression or the antidepressant
 - If existing SD, or very concerned about developing it, use a first-line antidepressant with a more favourable SD profile
 - Bupropion, agomelatine, vortioxetine, mirtazapine and moclobemide
 - For AISD where a change in treatment is not deemed reasonable, adding an 'antidote' is the most effective strategy. Also consider watchful waiting; scheduling sexual activity; dose reduction; drug holiday, and/or behavioural and/or complimentary intervention.
 - For AISD where a treatment change can be considered, switch to an agent with lower SD risk (as above)